Shoulder Injuries in the Industrial Athlete

New Frontiers: Transition Support and Achieving Mission Success
2009 NASA Occupational Health Meeting
Thursday July 16, 2009

Robert J. Dimeff, MD Director of PCSM Director of Education



Shoulder Injuries in the Industrial Athlete Objectives

- Locate important anatomic landmarks
- Know important history questions to assist in the evaluation of shoulder injuries
- Perform an appropriate shoulder exam
- Review common overuse and traumatic shoulder injuries
- Be able to treat common shoulder disorders

Shoulder Injuries in the Industrial Athlete

- Anatomy
- Biomechanics
- History
- Examination
- Common Injuries



Shoulder Injuries in the Industrial Athlete

- Anatomy
- Biomechanics
- History
- Examination
- Common Injuries




Shoulder Anatomy

- Bones
- Joints and Ligaments
- Muscles
- Neurovascular
- Bursae




Bones Clavicle

- S shaped
- Only bone to connect arm to axial skeleton
- Medial tubular/sternum
- Lateral flat/acromium
- Rotates along long axis
- Stabilize, protect plexus




Bones **Proximal Humerus**

- Spherical
- 130° inclination
- 30° retroversion
- 1/3 articular cartilage
- Greater and lesser tuberosities
- Intertubercular groove




Bones Scapula

- Body
- Spine
- Scapular neck
- Coracoid process
- Glenoid fossa
- Acromium




Bones Scapula

- Body
- Spine
- Scapular neck
- Coracoid process
- Glenoid fossa
- Acromium




Bones Scapula

- Body
- Spine
- Scapular neck
- Coracoid process
- Glenoid fossa
- Acromium




Joints and Ligaments Sternoclavicular joint

- Only joint to connect arm to axial skeleton
- Unstable ball and socket
- 1st rib cartilage
- 35° elevation
- 50° rotation




Joints and Ligaments Sternoclavicular joint

- Intraarticular disc
- Ligaments
 - ◆ costoclavicular
 - ◆ interclavicular
 - ◆ sternoclavicular




Joints and Ligaments Acromioclavicular joint

- Only connection of clavicle to scapula
- Hyaline becomes fibrocartilage with aging
- Intrarticular disc
- 40° of rotation
- Transmit forces from extremity to axis




Joints and Ligaments Acromioclavicular joint


- AC ligaments
 - ◆ AP stability
 - ◆ axial distraction
- Coracacoclavicular ligs
 - Trapezoid-anterior scapular rotation
 - Conoid-posterior scapular rotation



- Great mobility
- Poor stability
- 3mm superior/inferior translation
- 5mm anterior/posterior translation




- Glenoid
 - ◆ Peripheral cartilage
- Labrum
 - ◆ Fibrocartilage
 - Thick periphery
 - ◆ GH lig and biceps tendon




- GH capsule
 - ◆ Twice SA humeral head
 - ◆ Synovial lining
 - ◆ Minimal fluid
 - Negative intraarticular pressure
 - ◆ Check rein




- GH ligaments
 - ◆ Collagen thickening
 - **◆** Superior
 - ◆ Middle
 - ◆ Inferior
 - ◆ Static stability
 - Primary stabilizer at extremes of motion







- Coracohumeral ligament
 - ◆ Coracoid to greater tub
 - ◆ Suspension of humerus
 - Inferior instability



- Coracohumeral ligament
 - ◆ Coracoid to greater tub
 - ◆ Suspension of humerus
 - ◆ Inferior instability
- Coracoacromial ligament



- Coracohumeral ligament
 - ◆ Coracoid to greater tub
 - ◆ Suspension of humerus
 - ◆ Inferior instability
- Coracoacromial ligament
- Transverse humeral lig
 - Connect tuberosities
 - Maintain biceps tendon




Joints and Ligaments Scapulothoracic articulation

- Not a true joint
- Movement of scapula on chest wall
- Multiple muscles coordinate scapular movement




- Rotator cuff
 - Compresseshumeral head
 - Centers humerus
 - ◆ Active stabilizer
 - Resists displacing forces




Supraspinatus - abduct/


depress/compress




- Supraspinatus abduct/ depress/compress
- Infraspinatus ER/ab/ad
- Teres minor ER/ad




- Supraspinatus abduct/ depress/compress
- Infraspinatus ER/ab/ad
- Teres minor ER/ad
- Subscapularis -IR/ab/ad/flex/extend




- Biceps flex, supinate, depress humerus
- Coracobrachialis-flex/ad
- Pec major flex/ad/IR
- Pec minor draw scapula forward and downward



- Subclavius hold shoulder down and forward
- Deltoid primary mover
- Triceps ex/ad
- Teres major ad/ex/IR
- Serratus stabilize, rotate, draw scapula forward




- Rhomboids draw scapula medial, depress shoulder
- Levator scapula elevate and rotate scapula
- Lattisimus dorsi ex/ad/IR
- Trapezius raises, lowers, rotates, adducts scapula




Neurovascular

- Brachial Plexus
 - **♦** Roots
 - **◆**Trunks
 - ◆ Divisions
 - ◆ Cords
 - ◆ Nerves
- Axillary Artery
- Subclavian Vein




Neurovascular

- Brachial Plexus
 - **♦** Roots
 - ◆ Trunks
 - ◆ Divisions
 - ◆ Cords
 - ◆ Nerves
- Axillary Artery
- Subclavian Vein



Bursae

- Bursae
 - ◆ Subacromial
 - ◆ Subdeltoid
 - ◆ Subscapular
 - ◆ Biceps
 - ◆ Scapulothoracic



Shoulder Injuries in the Industrial Athlete

- Anatomy
- Biomechanics
- History
- Examination
- Common Injuries




Shoulder Biomechanics

- Result of 4 joints moving simultaneously and synchronously
- Humerus moves 2x as fast and far as scapula
- Abduction 180° -- Adduction 45°
- Flexion 160° -- Extension 45°
- Internal Rotation 90° -- External Rotation 100°
- Scapula -- elevate/depress, pro/retract


Shoulder Biomechanics

- Sternoclavicular joint
 - ♦ 35° anterior/posterior
 - ♦ 35° elevation
 - ♦ 45° rotation



Shoulder Biomechanics

- Sternoclavicular joint
 - ◆ 35° anterior/posterior
 - ♦ 35° elevation
 - ♦ 45° rotation
- Acromioclavicular joint
 - ◆ 20° rotation first 20°
 - ◆ 20° rotation last 40°




Shoulder Biomechanics

- Scapulothoracic joint
 - ◆ none during first 60°
 flexion or 30° abduction
 - synchronous with GH
 - ◆ 3° GH for each 2° ST




Shoulder Biomechanics

- Glenohumeral joint
 - ◆ 2/3 ab/flex, most ad/ext, all rotation
 - gliding and rolling
 - ◆ 3mm upward shift
 - instant center of rotation
 - effect of SC and AC joint and scapula motion
 - stability




Shoulder Biomechanics glenohumeral stability-static

- Bone
- Labrum
- GH Ligaments
- Concavity-compression
- Capsule
- Finite joint volume



Shoulder Biomechanics glenohumeral stability-dynamic

- Passive muscle bulk
- Muscle contraction
 - ◆ compresses joint surface
 - stiffens capsule
 - physical barrier
 - ◆ restraint
 - redirects joint reactive force to center of glenoid



Shoulder Injuries in the Industrial Athlete

- Anatomy
- Biomechanics
- History
- Examination
- Common Injuries





- History and exam essential for accurate DX
- Accurate diagnosis is required for successful treatment of any shoulder disorder
- Mechanism of injury is most important factor





- Chief complaint
 - ◆ Pain
 - ◆ Instability
 - ♦ Weakness
 - ◆ Loss of motion
 - Functional disability
 - Deformity
 - Catching/crepitus






- Symptom onset-acute or chronic
- Qualitate and quantitate symptoms
- Activities that aggravate/relieve symptoms
- Night, exertional, radiating pain
- Paresthesias, dead arm, weakness
- Looseness, slipping sensation, shoulder "popping out of joint"

- Muscle atrophy or fatigue
- Clicking, catching, grinding, grating
- Edema
- Coolness, cyanosis
- Trauma
- Overuse risk factors



- Age, occupation, avocation, and handedness
- Job demands
- General medical health
- Medication
- Connective tissue diseases
- Workers compensation
- Litigation



Shoulder Injuries in the Industrial Athlete

- Anatomy
- Biomechanics
- History
- **■** Examination
- Common Injuries



Examination

- Inspection-atrophy, swelling, discoloration
- ROM-symmetric, compensatory changes
- Palpation-deformity, tenderness
- Strength
- Impingement and stability
- Neck exam
- Neurovascular exam-sensation and reflexes

Examination range of motion

Examination palpation

Examination strength and impingement testing

Examination stability sitting

Examination stability supine

Examination radiographs

- AP, true AP
- Westpoint
- Axillary
- Outlet view
- Y view



Examination

radiographs

- AP, true AP
- Westpoint
- Axillary
- Outlet view
- Y view




Examination radiographs

- AP, true AP
- Westpoint
- Axillary
- Outlet view
- Y view





- AC/SC joints
- Scapula





Shoulder Injuries in the Industrial Athlete

- Anatomy
- Biomechanics
- History
- Examination
- Common Injuries



Common Shoulder Disorders

- SC dislocation
- Clavicle fracture
- AC sprain
- AC osteolysis
- Impingement
- Instability


- Subluxation and dislocation
- Labral tear
- Bicipital tendinitis
- Adhesive capsulitis
- Referred pain

Common Shoulder Disorders

- SC dislocation
- Clavicle fracture
- AC sprain
- AC osteolysis
- Impingement
- Instability

- Subluxation and dislocation
- Labral tear
- Bicipital tendinitis
- Adhesive capsulitis
- Referred pain


- Only articulation between arm and axial skeleton
- 95% anterior
- Anterior = indirect force to PL or AL shoulder
- Posterior = direct blow to medial clavicle




- Only articulation between arm and axial skeleton
- 95% anterior
- Anterior = indirect force to PL or AL shoulder
- Posterior = direct blow to medial clavicle



- Reduction
 - ◆ Anterior = Axial traction, downward pressure on clavicle
 - Posterior = Axial traction, upward pull
- Figure of 8, sling, ice, analgesics
- Injection



- Reduction
 - ◆ Anterior = Axial traction, downward pressure on clavicle
 - Posterior = Axial traction, upward pull
- Figure of 8, sling, ice, analgesics
- Injection



Common Shoulder Disorders

- SC dislocation
- Clavicle fracture
- AC sprain
- AC osteolysis
- Impingement
- Instability

- Subluxation and dislocation
- Labral tear
- Bicipital tendinitis
- Adhesive capsulitis
- Referred pain

Clavicle fracture

- Multiple classification systems
- 80% middle, 5% medial, 15% distal
- Fall on shoulder or outstretched arm, direct trauma
- Medial physis fuses at age 22-25



Clavicle fracture

- Figure of 8, shoulder immobilizer, exogen
- Displaced or distal may require operative fixation
- Rehab after period of immobilization
- Return to full activity in 4-12 weeks



Clavicle fracture

- Figure of 8, shoulder immobilizer, exogen
- Displaced or distal may require operative fixation
- Rehab after period of immobilization
- Return to full activity in 4-12 weeks




Common Shoulder Disorders

- SC dislocation
- Clavicle fracture
- AC sprain
- AC osteolysis
- Impingement
- Instability

- Subluxation and dislocation
- Labral tear
- Bicipital tendinitis
- Adhesive capsulitis
- Referred pain


AC sprain

- 6 types, I, II, III most common in sports
- AC and CC (trapezoid/conoid) ligaments



AC sprain

- 6 types, I, II, III most common in sports
- AC and CC (trapezoid/conoid) ligaments



AC sprain


- Fall on point of adducted shoulder or outstretched hand, direct trauma
- Sling, ice, analgesics, strap, injection, Kenny-Howard sling
- Surgery for IV, V, VI, and maybe III



position; and (3) the halter pulls both the shoulder and the sling inward.

AC joint osteolysis

- Weightlifters
- Localized shoulder pain
- Swelling, deformity, tenderness
- Impingement signs
- Rest, ice, analgesics
- Injection
- Surgical excision




- SC dislocation
- Clavicle fracture
- AC sprain
- AC osteolysis
- Impingement
- Instability

- Subluxation and dislocation
- Labral tear
- Bicipital tendinitis
- Adhesive capsulitis
- Referred pain


Impingement

- Beware of instability in younger athlete
- Positional pain,night pain
- Rest, ice, analgesics
- Rehab exercises



Impingement

- Kinetic chain
- Injections
- Modify activity
- Nerve entrapment
- MRI/msk US
- Beware RC tear
- May require surgery



Impingement

- Kinetic chain
- Injections
- Modify activity
- Nerve entrapment
- MRI/msk US
- Beware RC tear
- May require surgery



- SC dislocation
- Clavicle fracture
- AC sprain
- AC osteolysis
- Impingement
- Instability

- Subluxation and dislocation
- Labral tear
- Bicipital tendinitis
- Adhesive capsulitis
- Referred pain

Instability

- Anterior, posterior, inferior, MDI
- Pain during specific range of motion
- Dead arm
- Sulcus sign
- Apprehension




Instability

- Individualized treatment
 - severity of symptoms
 - * direction(s)
 - ◆ age and activity level
 - etiology
 - ◆ timing



Instability

- Rest, ice, analgesics
- Rehab exercises
- Posture
- Brace
- Traumatic often need surgery
- Atraumatic may require surgery



- SC dislocation
- Clavicle fracture
- AC sprain
- AC osteolysis
- Impingement
- Instability


- Subluxation and dislocation
- Labral tear
- Bicipital tendinitis
- Adhesive capsulitis
- Referred pain

Subluxation and dislocation


- Check NV status
- Relocation
 - axial traction and ER for anterior
 - axial traction and IR for posterior
- Sling for comfort
- ER brace *Itio*, et al











Subluxation and dislocation



Subluxation and dislocation

- Ice, analgesics
- Individualize treatment
- High recurrence rate < 20
- Rehabilitation exercises
- Brace
- Activity modification
- Often operate after first dislocation




- SC dislocation
- Clavicle fracture
- AC sprain
- AC osteolysis
- Impingement
- Instability

- Subluxation and dislocation
- Labral tear
- Bicipital tendinitis
- Adhesive capsulitis
- Referred pain

Positional pain, pop, instability


- Throwers + lifters
- Impingement signs
- O'Brien's
- MR arthrogram
- Conservative vsSurgical treatment

Type II



- Positional pain, pop, instability
- Throwers + lifters
- Impingement signs
- O'Brien's
- MR arthrogram
- Conservative vsSurgical treatment

Type IV




Positional pain, pop,

instability

■ Throwers + lifters

■ Impingement signs

- O'Brien's
- MR arthrogram
- Conservative vsSurgical treatment



- Positional pain, pop, instability
- Throwers + lifters
- Impingement signs
- O'Brien's
- MR arthrogram
- Conservative vsSurgical treatment








- SC dislocation
- Clavicle fracture
- AC sprain
- AC osteolysis
- Impingement
- Instability

- Subluxation and dislocation
- Labral tear
- Bicipital tendinitis
- Adhesive capsulitis
- Referred pain

Bicipital tendinitis

- Anterior pain with flexion and supination
- Instability, SLAP, impingement
- Tender bicipital groove, + speed



Bicipital tendinitis

- Treat as impingement
- Modalities
- Injection
- Pain free after rupture



- SC dislocation
- Clavicle fracture
- AC sprain
- AC osteolysis
- Impingement
- Instability

- Subluxation and dislocation
- Labral tear
- Bicipital tendinitis
- Adhesive capsulitis
- Referred pain


Adhesive capsulitis

- 40-60 year old females
- May be traumaticor atraumatic
- Pain, freeze, thaw
- Loss of active and passive motion




Adhesive capsulitis

- May have severe,mild, or no pain
- Moist heat
- Modalities
- Analgesics
- ROM exercises
- PRE



Adhesive capsulitis

- Cortisone injection in inflammatory phase
- High volume arthrogram
- Manipulation
- Educate patient
- Be patient




- SC dislocation
- Clavicle fracture
- AC sprain
- AC osteolysis
- Impingement
- Instability

- Subluxation and dislocation
- Labral tear
- Bicipital tendinitis
- Adhesive capsulitis
- Referred pain

Referred pain

- Cervical
- Upper extremity
- Brachial plexus
- Cardiac disease
- Pulmonary disease
- GI, diaphragmatic
- Splenic
- Malignancies



Thank You

